

# A hyperactive *piggyBac* transposase for mammalian applications

Kosuke Yusa<sup>a,1</sup>, Liqin Zhou<sup>b,1</sup>, Meng Amy Li<sup>a</sup>, Allan Bradley<sup>a,2</sup>, and Nancy L. Craig<sup>b,2</sup>

<sup>a</sup>Wellcome Trust Sanger Institute, Cambridge CB10 1SA, United Kingdom; and <sup>b</sup>Department of Molecular Biology and Genetics, The Howard Hughes Medical Institute, Johns Hopkins University School of Medicine, Baltimore, MD 21205-2185

Edited by Neal G. Copeland, Institute of Molecular and Cell Biology, Proteos, Singapore, and approved December 15, 2010 (received for review June 15, 2010)

DNA transposons have been widely used for transgenesis and insertional mutagenesis in various organisms. Among the transposons active in mammalian cells, the moth-derived transposon *piggyBac* is most promising with its highly efficient transposition, large cargo capacity, and precise repair of the donor site. Here we report the generation of a hyperactive *piggyBac* transposase. The active transposition of *piggyBac* in multiple organisms allowed us to screen a transposase mutant library in yeast for hyperactive mutants and then to test candidates in mouse ES cells. We isolated 18 hyperactive mutants in yeast, among which five were also hyperactive in mammalian cells. By combining all mutations, a total of 7 aa substitutions, into a single reading frame, we generated a unique hyperactive *piggyBac* transposase with 17-fold and ninefold increases in excision and integration, respectively. We showed its applicability by demonstrating an increased efficiency of generation of transgene-free mouse induced pluripotent stem cells. We also analyzed whether this hyperactive *piggyBac* transposase affects the genomic integrity of the host cells. The frequency of footprints left by the hyperactive *piggyBac* transposase was as low as WT transposase (~1%) and we found no evidence that the expression of the transposase affects genomic integrity. This hyperactive *piggyBac* transposase expands the utility of the *piggyBac* transposon for applications in mammalian genetics and gene therapy.

reprogramming | gene correction

DNA transposons are genetic elements that can mobilize from one location to another in the host genome. These have been used as laboratory tools for transgenesis and insertional mutagenesis in a wide range of model organisms such as *Drosophila* (1, 2), *Caenorhabditis elegans* (3, 4), and plants (5). However, their application to mammalian genetics had been hampered because of the lack of active transposons in mammals. Approximately a decade ago, the first active DNA transposon in mammals, *Sleeping Beauty*, was reconstructed from fossilized transposon sequences found in the salmonid genome (6). This pioneer work has greatly expanded the repertoire of tools for mammalian genetics. Germline transposition has accelerated the generation of mutant mice and rats (7–11), and somatic transposition has opened up numerous possibilities to conduct forward genetic screens in vivo such as cancer gene discovery in solid tumors (12–15). Furthermore, DNA transposons hold great promise for gene therapy as nonviral vehicles (16). Since the generation of the *Sleeping Beauty* transposon, a number of transposons from different families have been reported to show active transposition in mammalian cells. Among them, the *piggyBac* transposon isolated from cabbage looper moth *Trichoplusia ni* is most promising because of a variety of unique characteristics, namely exhibiting the most efficient transposition in mammalian cells, the ability of the transposase to form functional protein fusions, large cargo capacity, and traceless excision, i.e., its excision restores the donor site to its pretransposon state and leaves no trace of transposon insertion (17–20). Taking advantage of these unique characteristics, we have recently demonstrated the generation of factor-free mouse induced pluripotent stem (iPS) cells (21).

The DNA transposon system consists of two components: a DNA element flanked by two terminal inverted repeats (IRs) and a transposase that catalyzes the transposon's mobilization by a "cut-and-paste" mechanism. The transposases first bind to the IRs, then excise the DNA segment flanked by the IRs from the genome (i.e., cut) and finally reintegrate the segment into a new location (i.e., paste). Thus, engineering the transposase is central to increasing the transposition efficiency. This has been successfully applied to the *Sleeping Beauty* transposon system. The most recent version of the *Sleeping Beauty* transposase (SB100) shows a marked hyperactivity compared with the original transposase (22). We have previously demonstrated that a mammalian codon-optimized version of the *piggyBac* transposase (PBase) mediates more efficient transposition than the original insect version, a 20-fold increase in "plasmid-to-genome" transposition (20), and elevated rates of chromosomal transposition (23).

Here we report the generation of a hyperactive mutant of PBase. We first established a yeast assay to efficiently screen for mutations in the PBase, which give rise to hyperactivity. Individual candidate mutants were then verified for their transposition activity in mouse ES cells. Finally, we combined all the mutations into one sequence and generated a hyperactive PBase (hyPBase), which shows more than 10-fold higher rates of transposition than the WT mammalian codon-optimized PBase (mPBase).

## Results

**Isolation of hyPBase Mutants in *S. cerevisiae*.** We have previously reported that the insect *piggyBac* element can transpose in *Saccharomyces cerevisiae* and described assays for both transposon excision and integration (24). We have now used the excision assay to isolate hyperactive transposase mutants (Fig. 1A). In our two-plasmid *piggyBac* excision system in a yeast *URA3<sup>-</sup>* strain, the transposon donor plasmid carries a *URA3::actin* intron cassette (25) containing a 2.1-kb mini-*piggyBac* transposon in the actin intron. The actin intron::mini-*piggyBac* segment is too large to be spliced and thus the strain is a uracil auxotroph. Upon excision of the mini-*piggyBac* element, however, the intron can be spliced, reverting the strain to uracil prototrophy. Thus, measuring the frequency of reversion to ura<sup>+</sup> is a convenient assay for transposition. The transposase is supplied by a second plasmid containing the PBase gene under the galactose-inducible control of the *GALS* promoter (26).

We generated mutant pools of PBase DNA by error-prone PCR and introduced the pooled DNA into a yeast expression vector by

Author contributions: K.Y., L.Z., A.B., and N.L.C. designed research; K.Y., L.Z., and M.A.L. performed research; K.Y., L.Z., A.B., and N.L.C. analyzed data; and K.Y., L.Z., A.B., and N.L.C. wrote the paper.

The authors declare no conflict of interest.


This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

<sup>1</sup>K.Y. and L.Z. contributed equally to this work.

<sup>2</sup>To whom correspondence may be addressed. E-mail: ncraig@jhmi.edu or abradley@sanger.ac.uk.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1008322108/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1008322108/-DCSupplemental).




**Fig. 1.** Hyperactive mutant screening in yeast. (A) Schematic representation of the excision assay in yeast. The *URA3* gene is separated by the actin intron containing the mini-*piggyBac* transposon, which completely disturbs the normal splicing. After excision, the actin intron is spliced out normally and the *URA3* gene is restored. (B) Relative activities of hypBase mutants in yeast. These values are the median of assays of 10 colonies of WT and each mutant type performed in glucose, i.e., without induction of the GAL5 promoter. The absolute value of transposition promoted by WT transposase was  $4.7 \times 10^{-4}$  ura<sup>+</sup> cells/total cells.

homologous recombination. The mutant library was subsequently screened using the ura<sup>+</sup> reversion assay and the PBase mutants that gave rise to the higher number of ura<sup>+</sup> revertants than the WT PBase were isolated. In the initial screen, we examined 10,000 transformants and obtained approximately 200 candidates. The excision frequencies of these candidates were then analyzed quantitatively and we consequently isolated 18 hyperactive mutants, which showed two- to sevenfold increases in the excision activity (Fig. 1B).

**hypBase in Mammalian Cells.** We next investigated whether the PBase mutants isolated in yeast could also show elevated transposition in mammalian cells. The coding sequence of each mutant was transferred from the yeast expression vector to a CMV promoter-based mammalian expression vector. To find bona fide hyperactive mutants, we measured both excision and integration frequencies in mouse ES cells as illustrated in Fig. 2A and B, respectively. In the excision assay (Fig. 2A), we used an ES cell line that has a *piggyBac* transposon targeted into intron 2 of the *Hprt* gene (Fig. S14). The transposon carries a splice acceptor element, which disrupts the expression of the *Hprt* gene. The targeted ES cells are thus sensitive to HAT. When the transposon jumps out from the donor site, *Hprt* expression is restored and ES cells become resistant to HAT. In the integration assay (Fig. 2B), we cotransfected WT ES cells with the transposase expression vector and a transposon carrying a gene-trap cassette. When the transposon integrates into genes expressed in ES cells, the puromycin resistant gene is expressed; thereby ES cells become resistant to puromycin.

The activity of the 18 candidate mutants and the WT PBase are shown in Fig. 2C. Contrary to the results in the yeast assay, 13 mutants showed similar or weaker activity in ES cells in both excision and integration compared with the WT PBase. For instance, the second-strongest mutant in yeast (Q591P) did not show any significant increase in transposition in ES cells. The fourth strongest mutant in yeast (M194V) had very low activity in ES cells. Nevertheless, five mutants, namely I30V/G165S, S103P, M282V,




**Fig. 2.** Comparison of the transposase mutants and generation of hyperactive transposase in mouse ES cells. (A) Schematic representation of the excision assay in ES cells. The *Hprt* gene is disrupted by the *piggyBac* transposon carrying a gene-trap unit; thus, the ES cells are sensitive to HAT. When the transposon jumps out, the *Hprt* gene is restored, making cells resistant to HAT. (B) Schematic representation of the integration assay in ES cells. The *piggyBac* transposon vector carrying a gene-trap cassette was cotransfected into WT ES cells together with a transposase expression vector. When the transposon jumps into an active gene, the puromycin resistant gene is expressed; thus cells become resistant to puromycin. White boxes, exons; SA, splice acceptor site; IRES, internal ribosome entry site; *puΔtk*, the puromycin-resistant gene fused with the herpes simplex virus *thymidine kinase* gene; pA, poly-adenylation signal. (C) Relative excision and integration activities of the hyperactive mutant in ES cells (Upper) and protein expression of HA-tagged PBase and mutants in 293T cells (Lower). As a control, a WT transposase with the original insect-derived coding sequence was used. The mutants indicated by arrows were combined for the generation of the hyperactive transposase. Representative data are shown. Asterisk marks nonspecific band. (D and E) Comparison of hypBase with the WT transposase (mPBBase) in excision (D) and integration (E) assays. Data are shown as mean  $\pm$  SD ( $n = 3$ ). (F) Western blot analysis showing expression of HA-tagged mPBBase and hypBase in ES cells. Dilution factors are shown on the top of gel picture.  $\beta$ -Actin was used for loading controls. Asterisk marks nonspecific band.

S509G/N570S, and N538K, showed more than a twofold increase in activity in both the excision and integration assays. The fold changes of these mutants were similar between excision and integration, suggesting that the mutations mainly enhanced the excision reaction and the integration reaction remained unaffected. These outcomes could be caused by altered levels of protein expression. We assessed this by using Western blot analysis of HA-tagged mutant Pbases (Fig. 2C, Lower). Decreased transposition activities in the M194V, R281G, and G316E mutants were associated with marked reduction of their protein levels. The strongest mutant in both yeast and ES cells (M282V) had a slightly increased level of protein. The rest of the mutants showed similar protein levels to the WT PBase. Steady-state protein levels were thus correlated with the transposition activity in some, but not all, mutants. The mutants with unchanged protein levels most likely affected the mechanism and/or kinetics of transposition.

We then investigated whether combinations of these mutations can synergistically enhance the transposition efficiency.





**Fig. 4.** Improved generation of transgene-free iPSC cells using the hyPBBase. (A) A representative image of alkaline phosphatase staining of iPSC cell colonies generated using WT (Left) and hyperactive (Right) PBBase. (B) Numbers of iPSC cell colonies obtained from transfection of MEFs with 100 ng transposon and 100 ng transposase in a 12-well plate. Data are shown as mean  $\pm$  SD ( $n = 3$ ) (C) Number of transgene-free iPSC cell colonies generated by mPBBase or hyPBBase. Both primary iPSC cell lines (iPS25 and iPS28) have two transposon integrations. Representative data are shown. (D) PCR analysis showing transposon removal and no evidence of random integration of plasmids. (E) Precise repair of the excised site. All clones examined possess intact genomic sequences. The transposon donor sites are highlighted in red.

creased eightfold by using hyPBBase compared with the mPBBase, indicating that the hyPBBase enhanced reprogramming efficiency.

To generate factor-free iPSC cells, the integrated transposons need to be excised from the iPSC cell genome. In a previous study (21), the *piggyBac* excision efficiency was approximately  $1 \times 10^{-5}$  per cell, which was three orders of magnitude lower than the excision frequency at the *Hprt* locus. To test if hyPBBase could be a useful tool for transposon removal from iPSC cells, we transfected two primary iPSC cell lines (iPS25 and iPS28) (21) with mPBBase and hyPBBase and subjected the transfected cells to FIAU selection. As shown in Fig. 4C, hyPBBase removed the transposons approximately 20 times more efficiently than mPBBase. PCR analyses showed that all clones analyzed had lost the transposon and did not have random integrations of the PBBase expression vector (Fig. 4D). Importantly, we did not detect any footprint mutations at the primary transposon integration sites in both mPBBase- and hyPBBase-mediated excisions, showing the generation of transgene-free iPSC cells by the hyPBBase (Fig. 4E and F).

**PBBase Does Not Affect Genomic Integrity.** Transposases catalyze transposon excision from the host genome by introducing double-stranded breaks (DSBs) at both ends of the transposon. In principle, expression of the transposase may cause genomic instability as a result of DSBs at cryptic sites in the genome. To investigate this possibility, we conducted comparative genomic hybridization (CGH) on genomic DNA isolated from PBBase-exposed cells by using an Agilent 244K mouse genome array. We first compared two primary iPSC cell lines (iPS25 and iPS28; both carry two copies of the transposon; ref. 21) with 10 transposon-free iPSC cell lines derived from them following mPBBase- or hyPBBase-mediated excision. The results, summarized in Table S2, confirmed that the primary iPSC cell lines do not have any abnormalities compared with the donor MEFs. Half the transposon-free iPSC cell lines did not show any aberrations within the limits of CGH resolution, indicating that they possess unaltered genomes. The remaining half the cell lines, however, had deletions and/or amplifications ranging from 30 to 300 kb and one line, 25 $\Delta\Delta$ 1, had a large rearrangement on chromosome 8. None of the changes were linked to the original transposon integration sites. These genomic alterations could be caused by continuous transposition before trans-

poson loss or the transposase directly, or arise spontaneously during culture. As donor site deletions caused by transposon excision accounted for just 0.8% of total excision events and their sizes were less than 5 kb (Fig. 3), the genomic alterations found in transposon-free iPSC cell lines do not seem to be mediated by transposon excision. We assessed this further by isolating 15 transposon-free iPSC cell lines using hyPBBase from the iPS25 primary line and we compared these with 25 subclones from the same primary line using CGH analysis. Genomic alterations were found in 20% of transposon-free iPSC cell lines (3 of 15 lines) and 20% of subclones (5 of 25 lines; Table 1). Therefore, we did not find a statistically significant difference in genomic abnormality in hyPBBase-treated versus untreated iPSC cells (iPS25-hyPBBase, four of 17; iPS25-subclone, 5 of 25;  $P = 0.75$ , Fisher exact test).

We carried out two additional assays to investigate if PBBase itself can induce significant damage to the host genome. In the first assay, we investigated *Hprt* mutation frequency upon transposase exposure. WT ES cells were transfected with iPBase, mPBBase, or hyPBBase expression vectors ( $\sim 90\%$  transfection efficiency), cultured for 6 d, and subsequently subjected to 6TG selection. We observed a few *Hprt*-deficient colonies from iPBase and mPBBase-transfected cells, but none in the control (GFP-transfected) or hyPBBase-transfected cells (Table S3). The colonies that did appear occurred at background rates but none appeared following treatment with hyPBBase.

In the second assay, we analyzed DSB levels after transient expression of PBBase by using two surrogates, H2AX phosphorylation and foci as well as sister chromatid exchange (SCE). In these experiments, we expressed PBBase by mRNA transfection, which also can induce efficient transposition (Fig. S3A and B). First we analyzed H2AX phosphorylation, a biomarker of DNA damage (27), by Western blotting. All types of PBBase-transfected cells showed similar levels of H2AX phosphorylation compared with GFP-transfected and nontreated cells (Fig. S3C).  $\gamma$ H2AX foci in transfected NIH 3T3 analyzed by immunofluorescence were unaffected by PBBase expression (Fig. S3D and E). Last, we measured SCEs in PBBase-transfected ES cells. SCE is used as a cytogenetic biomarker to examine mutagenicity of chemical agents (28). SCEs are also induced by endonucleases (29). We did not observe any differences between nontreated and PBBase-

**Table 1. CGH analysis of iPS cell lines**

iPS cell line	Transposase	Aberration	Aberration type	Chr.	Size, kb	Found in other lines	No. of genes
iPS cell lines subcloned from iP525 (five abnormal lines of 25 lines analyzed)							
25-sub6	NA	Yes	1-copy deletion	4	58.3	No	2
25-sub8	NA	Yes	1-copy gain	5	524.8	No	8
			1-copy deletion	15	32.9	No	0
25-sub9	NA	Yes	1-copy deletion	6	1,847.6	Yes	0
25-sub16	NA	Yes	1-copy deletion	6	1,847.6	Yes	0
25-sub18	NA	Yes	1-copy deletion	5	92.8	No	0
Transposon-free iPS cell lines derived from iP525 (three abnormal lines of 15 lines analyzed)							
25-hy4	hyPBase	Yes	1-copy deletion	4	64.0	No	2
25-hy7	hyPBase	Yes	1-copy deletion	9	105.9	No	2
25-hy12	hyPBase	Yes	1-copy deletion	10	73.3	No	4

treated ES cells (Fig. S3 F and G). These results suggest that PBase itself does not seem to induce genomic instability.

Taken together, genomic alterations found in transposon-free iPS cells were highly likely to be acquired during culture and were not caused by expression of transposases.

### Discussion

In this study, we developed a hyperactive version of the *piggyBac* transposase and demonstrated its use to improve the efficiency of generation of the transgene-free iPS cells. We have also investigated in detail whether the use of *piggyBac* transposon and transposase cause genomic instability.

Two lines of evidence suggest that *piggyBac* transposition may be a host factor-independent reaction: first, its transposition is highly efficient in a wide range of organisms such as yeast (24), insects (30), planarian (31), the malaria parasite (32), and mammals (17); second, transposition can be reconstituted in vitro by using purified PBase and DNA elements (24). These characteristics of the *piggyBac* transposon allowed us to first screen a large number of transposase mutants in yeast and then test the candidates in mammals. Based on this strategy, we found five hyperactive mutants in ES cells among 10,000 mutants screened initially in yeast. As our mutant library in yeast was not saturated, more hyperactive mutants might be identified with this screening system in the future.

Despite the host-factor independence of *piggyBac* transposition, three fourths of the hyperactive mutants in yeast did not show hyperactivity in ES cells. There may be two factors contributing to this observation. First, species-specific factors may modulate the transposition positively or negatively. For example, the *Sleeping Beauty* transposase directly interacts with Miz-1 transcription factor (33) and HMGB1 (34), leading to the enhancement of transposition. Alternatively, epigenetic modifications may influence transposition frequency. The *Sleeping Beauty* transposase has an affinity for heterochromatic regions; hence, repressive epigenetic modification of the *Sleeping Beauty* transposon can increase the transposition frequency (35, 36). Although no transposase–host factor interaction in the *piggyBac* transposon system has been reported, if they do occur, PBase mutations might influence these interactions in a species-specific manner. Another possibility is that the reaction temperature affects catalytic activity and/or protein stability as mammalian cells are grown at very different temperatures from yeast. The activity of a yeast protein, FLP recombinase, was improved for mammalian use by increasing its stability at higher temperatures (37). Although the mechanism of how each mutation modulates transposition frequency remains to be determined, the mutants with different transposition efficiency in yeast and mammalian cell lines could be useful tools for further investigation of transposon biology.

DNA transposons are very useful as nonviral vehicles for genome engineering, which may be used for gene therapy. However,

safety concerns, especially genotoxicity, must be fully assessed. We have measured two aspects of the *piggyBac* transposon system: excision-induced genomic alterations at the donor site and a genome-wide assessment of transposase-induced genomic instability. Excision-induced genomic alterations are detected in 0.8% of excision events by mPBase and hyPBase transposases, which is much lower than the previously reported footprint frequency (~5–10%) (19). This difference might reflect the excision loci assayed or the number of excision events analyzed. Among excision-induced mutations, we found genomic alterations up to 5 kb from the donor site at very low frequency (~0.2% of excision events). These types of excision-induced mutations have not been described in previous reports (19). It is well recognized that mobilization of other types of DNA transposons can cause larger genomic rearrangements including insertion, deletion, duplication, inversion, and translocations, particularly by the *P*-element (38) and *Ac/Ds* elements (39–41) in *Drosophila* and *Maize*, respectively. In mice, transposition of *Sleeping Beauty* from a concatemer donor site can cause deletions and/or inversions of megabases of sequence as well as translocations (42). We did not observe such large and complex alterations following *piggyBac* excision. Although our results do not rule out the possibility of larger genomic alterations occurring, in general *piggyBac* excision from a single-copy donor site does not significantly affect genomic integrity.

A second concern is whether the *piggyBac* transposase itself causes genomic instability at sites other than a donor locus. To address this issue, we performed CGH analysis and compared 17 transposon-free iPS cells generated by hyPBase transfection and 25 subclones not exposed to hyPBase. Although we found that several cell lines carried deletions or amplifications, there were no obvious differences between these two groups in the nature of the detectable genomic aberrations such as their size, gain or loss of sequence, and the frequency with which they occurred. Moreover, we previously reported that mouse ES cells are genetically heterogeneous as a result of copy number change, with 30% to 40% of single cell subclones exhibiting copy number change (43). These results strongly suggest that the genomic alterations in the transposon-free iPS cells arose spontaneously and were not caused by PBase expression; hence, the hyPBase can mediate more efficient transposition without compromising the genomic integrity. Nevertheless, it might be worth noting that transposons can jump multiple times before they are integrated or lost and thus there is a small probability of footprint mutations. These mutations would be too small to be detected by currently available genome-wide methods. Whole-genome sequencing would have the resolution to detect such changes, and this might be considered before clinical use of iPS cells. Although we believe footprint mutations occur rarely, the development of transposases that can catalyze only excision but not integration or small compounds that can inhibit

only integration reaction might eliminate the chance of generating footprint mutations in applications of transposon removal.

The *piggyBac* transposase is amenable to modification; for instance, terminal fusion with activity modulator domains such as ERT2 and Gal4 DNA binding domains have been reported (18, 20, 44), and in this study we have used HA-tagged versions. These modifications allow us to regulate transposition in a spatiotemporal manner. In vivo applications of such engineered transposases are useful to address a variety of biological questions. This hyperactive *piggyBac* transposon system will further expand the use of transposons as tools for genome engineering such as insertional mutagenesis and gene therapy.

## Materials and Methods

**Plasmid Construction.** To construct mammalian expression vectors of the mutant *piggyBac* transposase, the BamHI-XhoI fragments containing the

mutant sequences in the yeast expression vector pGALS were transferred into the BamHI-XhoI site of pcDNA3 (Invitrogen). The point mutations were introduced into pCMV-mPBase (20) by site-directed mutagenesis to generate hyperactive transposases. The hyPBBase expression vector (pCMV-hyPBBase) is available upon request to the Sanger Institute Archives (<http://www.sanger.ac.uk/technology/clonerequests/>).

Construction of the targeting vectors, mutant screening in yeast, ES cell experiment, iPSC cell reprogramming, transposon excision in iPSC cells, and CGH analysis are described in *SI Materials and Methods*. Primer sequences are shown in Table S4.

**ACKNOWLEDGMENTS.** We thank J. Cooper and F. Law for technical assistance, R. Rad for comments on the manuscript, and the microarray team for data generation and analysis. K.Y. is funded by a postdoctoral fellowship of the Japan Society for the Promotion of Science. N.L.C. is an Investigator of The Howard Hughes Medical Institute. This work was supported by Wellcome Trust Grant WT077187.

- Bellen HJ, et al. (1989) P-element-mediated enhancer detection: A versatile method to study development in *Drosophila*. *Genes Dev* 3:1288–1300.
- Spradling AC, et al. (1995) Gene disruptions using P transposable elements: An integral component of the *Drosophila* genome project. *Proc Natl Acad Sci USA* 92: 10824–10830.
- Greenwald I (1985) *lin-12*, a nematode homeotic gene, is homologous to a set of mammalian proteins that includes epidermal growth factor. *Cell* 43:583–590.
- Moerman DG, Benian GM, Waterston RH (1986) Molecular cloning of the muscle gene *unc-22* in *Caenorhabditis elegans* by Tc1 transposon tagging. *Proc Natl Acad Sci USA* 83:2579–2583.
- Osborne BI, Baker B (1995) Movers and shakers: maize transposons as tools for analyzing other plant genomes. *Curr Opin Cell Biol* 7:406–413.
- Ivics Z, Hackett PB, Plasterk RH, Izsvák Z (1997) Molecular reconstruction of Sleeping Beauty, a Tc1-like transposon from fish, and its transposition in human cells. *Cell* 91: 501–510.
- Horie K, et al. (2003) Characterization of Sleeping Beauty transposition and its application to genetic screening in mice. *Mol Cell Biol* 23:9189–9207.
- Keng VW, et al. (2005) Region-specific saturation germline mutagenesis in mice using the Sleeping Beauty transposon system. *Nat Methods* 2:763–769.
- Kitada K, et al. (2007) Transposon-tagged mutagenesis in the rat. *Nat Methods* 4: 131–133.
- Izsvák Z, et al. (2010) Generating knockout rats by transposon mutagenesis in spermatogonial stem cells. *Nat Methods* 7:443–445.
- Carlson CM, et al. (2003) Transposon mutagenesis of the mouse germline. *Genetics* 165:243–256.
- Collier LS, Carlson CM, Ravimohan S, Dupuy AJ, Largaespada DA (2005) Cancer gene discovery in solid tumours using transposon-based somatic mutagenesis in the mouse. *Nature* 436:272–276.
- Dupuy AJ, Akagi K, Largaespada DA, Copeland NG, Jenkins NA (2005) Mammalian mutagenesis using a highly mobile somatic Sleeping Beauty transposon system. *Nature* 436:221–226.
- Keng VW, et al. (2009) A conditional transposon-based insertional mutagenesis screen for genes associated with mouse hepatocellular carcinoma. *Nat Biotechnol* 27: 264–274.
- Starr TK, et al. (2009) A transposon-based genetic screen in mice identifies genes altered in colorectal cancer. *Science* 323:1747–1750.
- Hackett PB, Largaespada DA, Cooper LJ (2010) A transposon and transposase system for human application. *Mol Ther* 18:674–683.
- Ding S, et al. (2005) Efficient transposition of the piggyBac (PB) transposon in mammalian cells and mice. *Cell* 122:473–483.
- Wu SC, et al. (2006) piggyBac is a flexible and highly active transposon as compared to sleeping beauty, Tol2, and Mos1 in mammalian cells. *Proc Natl Acad Sci USA* 103: 15008–15013.
- Wang W, et al. (2008) Chromosomal transposition of PiggyBac in mouse embryonic stem cells. *Proc Natl Acad Sci USA* 105:9290–9295.
- Cadiñanos J, Bradley A (2007) Generation of an inducible and optimized piggyBac transposon system. *Nucleic Acids Res* 35:e87.
- Yusa K, Rad R, Takeda J, Bradley A (2009) Generation of transgene-free induced pluripotent mouse stem cells by the piggyBac transposon. *Nat Methods* 6:363–369.
- Mátés L, et al. (2009) Molecular evolution of a novel hyperactive Sleeping Beauty transposase enables robust stable gene transfer in vertebrates. *Nat Genet* 41: 753–761.
- Liang Q, Kong J, Stalker J, Bradley A (2009) Chromosomal mobilization and reintegration of Sleeping Beauty and PiggyBac transposons. *Genesis* 47:404–408.
- Mitra R, Fain-Thornton J, Craig NL (2008) piggyBac can bypass DNA synthesis during cut and paste transposition. *EMBO J* 27:1097–1109.
- Yu X, Gabriel A (1999) Patching broken chromosomes with extranuclear cellular DNA. *Mol Cell* 4:873–881.
- Mumberg D, Müller R, Funk M (1994) Regulatable promoters of *Saccharomyces cerevisiae*: comparison of transcriptional activity and their use for heterologous expression. *Nucleic Acids Res* 22:5767–5768.
- Bonner WM, et al. (2008) GammaH2AX and cancer. *Nat Rev Cancer* 8:957–967.
- Perry P, Evans HJ (1975) Cytological detection of mutagen-carcinogen exposure by sister chromatid exchange. *Nature* 258:121–125.
- Obe G, Schunck C, Johannes C (1994) Induction of sister-chromatid exchanges by AluI, DNase I, benzonuclease and bleomycin in Chinese hamster ovary (CHO) cells. *Mutat Res* 307:315–321.
- Handler AM (2002) Use of the piggyBac transposon for germ-line transformation of insects. *Insect Biochem Mol Biol* 32:1211–1220.
- González-Estévez C, Momose T, Gehring WJ, Saló E (2003) Transgenic planarian lines obtained by electroporation using transposon-derived vectors and an eye-specific GFP marker. *Proc Natl Acad Sci USA* 100:14046–14051.
- Balu B, Shoue DA, Fraser MJ, Jr., Adams JH (2005) High-efficiency transformation of *Plasmodium falciparum* by the lepidopteran transposable element piggyBac. *Proc Natl Acad Sci USA* 102:16391–16396.
- Walisko O, et al. (2006) Sleeping Beauty transposase modulates cell-cycle progression through interaction with Miz-1. *Proc Natl Acad Sci USA* 103:4062–4067.
- Zayed H, Izsvák Z, Khare D, Heinemann U, Ivics Z (2003) The DNA-bending protein HMG1 is a cellular cofactor of Sleeping Beauty transposition. *Nucleic Acids Res* 31: 2313–2322.
- Yusa K, Takeda J, Horie K (2004) Enhancement of Sleeping Beauty transposition by CpG methylation: Possible role of heterochromatin formation. *Mol Cell Biol* 24: 4004–4018.
- Ikeda R, et al. (2007) Sleeping beauty transposase has an affinity for heterochromatin conformation. *Mol Cell Biol* 27:1665–1676.
- Buchholz F, Angrand PO, Stewart AF (1998) Improved properties of FLP recombinase evolved by cycling mutagenesis. *Nat Biotechnol* 16:657–662.
- Gray YH (2000) It takes two transposons to tango: Transposable-element-mediated chromosomal rearrangements. *Trends Genet* 16:461–468.
- Zhang J, Peterson T (1999) Genome rearrangements by nonlinear transposons in maize. *Genetics* 153:1403–1410.
- Zhang J, Peterson T (2004) Transposition of reversed Ac element ends generates chromosome rearrangements in maize. *Genetics* 167:1929–1937.
- Zhang J, Peterson T (2005) A segmental deletion series generated by sister-chromatid transposition of Ac transposable elements in maize. *Genetics* 171:333–344.
- Geurts AM, et al. (2006) Gene mutations and genomic rearrangements in the mouse as a result of transposon mobilization from chromosomal concatemers. *PLoS Genet* 2: e156.
- Liang Q, Conte N, Skarnes WC, Bradley A (2008) Extensive genomic copy number variation in embryonic stem cells. *Proc Natl Acad Sci USA* 105:17453–17456.
- Lacoste A, Berenshteyn F, Brivanlou AH (2009) An efficient and reversible transposable system for gene delivery and lineage-specific differentiation in human embryonic stem cells. *Cell Stem Cell* 5:332–342.